

Does Iron Supplementation Improve Performance in Iron-Deficient Nonanemic Athletes?

Amity Rubeor, DO, CAQSM,^{*,†} Carmen Goojha, MD,[‡] Jeffrey Manning, MD, CAQSM,[†] and Jordan White, MD, MPH[†]

Context: Supplementing iron-deficient nonanemic (IDNA) athletes with iron to improve performance is a trend in endurance sports.

Objectives: To investigate the benefits of iron on performance, identify a ferritin level cutoff in IDNA athletes, and determine which iron supplementation regimens are most effective.

Data Sources: A search of the PubMed, CINAHL, Embase, ERIC, and Cochrane databases was performed in 2014 including all articles. Citations of pertinent review articles were also searched. In 2017, the search was repeated.

Study Selection: Inclusion criteria comprised studies of level 1 to 3 evidence, written in the English language, that researched iron supplementation in nonanemic athletes and reported performance outcomes.

Study Design: Systematic review.

Level of Evidence: Level 3.

Data Extraction: The search terms used included *athletic performance, resistance training, athletes, physical endurance, iron, iron deficiency, supplement, non-anemic, low ferritin, ferritin, ferritin blood level, athletes, and sports.*

Results: A total of 1884 studies were identified through the initial database search, and 13 were identified through searching references of relevant review articles. A subsequent database search identified 46 studies. Following exclusions, 12 studies with a total of 283 participants were included. Supplementing IDNA athletes with iron improved performance in 6 studies (146 participants) and did not improve performance in the other 6 studies (137 participants). In the 6 studies that showed improved performance with iron supplementation, all used a ferritin level cutoff of ≤ 20 $\mu\text{g/L}$ for treatment. Additionally, all studies that showed improved performance used oral iron as a supplement.

Conclusion: The evidence is equivocal as to whether iron supplementation in IDNA athletes improves athletic performance. Supplementing athletes with ferritin levels < 20 $\mu\text{g/L}$ may be more beneficial than supplementing athletes with higher baseline ferritin levels.

Keywords: medical aspects of sports; athletic training; iron supplementation; performance

Iron is an essential mineral necessary for delivering oxygen to tissue throughout the body as well as serving important roles in metabolism, respiration, and immune function. The body maintains stores of iron and carefully sustains a balance between iron lost, iron absorbed, and iron stored.^{10,21-23,38}

Ferritin is the cellular storage protein for iron and is generally reflective of total body iron stores. Therefore, serum ferritin (sFer) is the measure most commonly used to predict total body iron stores. Notably, ferritin also has a role as an acute-phase

reactant, and ferritin levels measured in individuals with acute or chronic illness resulting in systemic inflammation will not appropriately predict iron stores.^{10,21,48}

Iron deficiency anemia (IDA) is a significant health concern worldwide and results in a host of symptoms, including weakness and fatigue. IDA represents the third, final, and most severe stage of iron deficiency.^{21,38} The 2 initial stages, which are precursors to IDA, can be collectively referred to as iron deficiency nonanemia (IDNA). The first and least severe stage of

From [†]Department of Family Medicine, Warren Alpert Medical School of Brown University, Providence, Rhode Island, and [‡]Family Physician, Sechelt, British Columbia, Canada
*Address correspondence to Amity Rubeor, DO, CAQSM, Family Health and Sports Medicine, 65 Sockanosset Cross Road, Suite 301, Cranston, RI 02920 (email: amity_rubeor@brown.edu).

The authors report no potential conflicts of interest in the development and publication of this article.

DOI: 10.1177/1941738118777488

© 2018 The Author(s)

iron deficiency (nonanemia) is marked by a fall in sFer resulting from the depletion of total body iron stores while maintaining normal levels of other iron indices and normal hemoglobin. The second stage (also nonanemia) is marked by not only low sFer but also low serum iron or decreased transferrin saturation and increased total iron binding capacity (TIBC).^{21,38} Once iron stores and transport iron have been sufficiently depleted, the body can no longer keep up with the demands of hemoglobin synthesis, and the third and final stage (IDA) results.^{21,23,38}

Monitoring sFer concentration has become common practice in endurance athletes at all levels, and not just athletes with a history of IDA. There is widespread belief that low ferritin, even in the context of normal hemoglobin levels (>12 g/dL), is the culprit for fatigue and poor performance.³⁸ Athletes with IDNA routinely use iron supplementation regimens in an effort to improve performance.^{5,10,21,38,48} One significant limitation in the current body of literature concerning IDNA is a lack of standardization for what is considered a “low” ferritin level. Although many studies and most laboratories generally define low ferritin in the range of 12 to 23 µg/L,¹⁰ the ferritin ranges used in research are not standardized, and due to the lack of clinical consensus on what constitutes “low” ferritin, published studies use a variety of different values to represent low normal levels, ranging from 12 to 40 µg/L.^{4,11,19,20,27-29,35,36,39,47}

There is mixed evidence regarding the potential negative impact of IDNA on endurance performance in athletes. There are no known negative effects of IDNA on the oxygen-carrying capacity of athletes.⁵ The hypothesized mechanism for potentially decreased endurance in IDNA athletes is decreased extraction and utilization of oxygen bound to hemoglobin due to the inactivity of iron-dependent oxidative enzymes and respiratory proteins.^{5,10} Thus, supplementation with iron could theoretically correct this deficit and improve performance. The 2015 meta-analysis by Burden et al⁵ supported this hypothesis and concluded that iron supplementation does indeed improve the aerobic capacity of IDNA athletes, measured by maximal VO₂.⁵

Although that meta-analysis⁵ found that iron supplementation can improve aerobic capacity, there is no consensus on the optimal effective route, dose, or frequency of iron supplementation in athletes. Burden et al⁵ included studies with intramuscular and oral iron supplementation as well as dosing regimens, which varied from 1 to 3 times per day at doses ranging from 10 to 209 mg.⁵ In addition, the meta-analysis did not distinguish between the 2 stages of IDNA.⁵

The purpose of this study was to conduct a systematic review of iron supplementation in IDNA athletes.

METHODS

Literature Search

The PubMed, CINAHL, Embase, ERIC, and Cochrane databases were systematically searched using the following search terms: *athletic performance, resistance training, athletes, physical endurance, iron, iron dietary, ferritin, supplement, non-anemic,*

low ferritin, and iron storage. Our preliminary search eliminated animal studies and studies in languages other than English. Additional studies were identified on reviewing references of pertinent review articles. Prior to publication, we performed an additional search to determine whether there had been published studies focusing on a ferritin cutoff of ≤20 µg/L for supplementation. We searched Embase and PubMed using the following terms: *ferritin, ferritin blood level, athletes, and sports.*

Inclusion and Exclusion Criteria

Studies were further screened according to predefined inclusion and exclusion criteria. Inclusion criteria consisted of: (1) randomized controlled trials; (2) athlete participants (recreational, club, high school, college, amateur, or professional athletes); (3) participants defined by the study authors as having IDNA or low ferritin without anemia; (4) at least 2 study groups, 1 of which received iron supplementation (dietary, intravenous, intramuscular, or pill form) as an intervention; and (5) outcomes comprising at least 1 measure of performance. Exclusion criteria included: (1) studies in which performance was measured subjectively; (2) studies including participants both with and without iron deficiency; (3) participants who could not be confirmed as athletes, for example, military recruits in basic combat training; and (4) studies that did not use a placebo as control. Three reviewers participated in screening and thoroughly reviewing each eligible study. Disagreements were resolved by consensus (Figure 1).

RESULTS

Overall, 12 total studies were included in our systematic review. In these 12 studies, there were a total of 283 participants (age range, 13–41 years; 257 females, 26 males). A summary of these studies is presented in Table A1 in the Appendix (available in the online version of this article).

Performance Measures

The studies included in this review used a wide variety of performance measures. Treating IDNA athletes with iron supplementation improved performance in 6 studies (146 participants); however, the other 6 studies (137 participants) did not show any improvement (Table A2 in the Appendix).

Ferritin Parameters

There was inconsistency across studies regarding a ferritin level diagnostic for IDNA. The studies included used ferritin cutoffs >20 or <20 µg/L (see Figure 2 and Table A3 in the Appendix).

Iron Supplementation Protocols

Of the 6 studies that found improved performance after iron supplementation, a variety of iron treatment protocols with differing preparations, doses, frequency, and lengths of treatment were used (see Figure 3 and Tables A4 and A5 in the Appendix).^{11,20,27,29,39,47}




Figure 1. Flow diagram of literature selection on the effects of iron supplementation in iron-deficient nonanemic (IDNA) athletes. RCT, randomized controlled trial.

DISCUSSION

Performance Improvement

The results of our review are equivocal and do not support the conclusion that iron supplementation *consistently* improves performance in endurance IDNA athletes. Studies included in this review used a variety of clinically applicable measures of performance, such as a shuttle run time, 3000-m

race pace, anaerobic speed test, maximal treadmill velocity, and treadmill endurance times.^{4,33,47} Given that each sport has its own physical demands, it is logical that optimal performance will be measured differently between sports. For example, although VO_{2max} is an excellent indicator of cardiorespiratory endurance, Garza et al²¹ theorized in 1997 that time to exhaustion “simulated field experience most closely.”




Figure 2. Iron formulations—performance improvement.




Figure 3. Iron formulations—no performance improvement.

Our conclusion that iron supplementation does not consistently improve performance in IDNA athletes differs from that of Burden et al,⁵ who concluded that iron supplementation improves aerobic capacity (measured by maximal VO_2) in IDNA athletes. However, that meta-analysis only included studies that used $\text{VO}_{2\text{max}}$ as the performance measure. On the other hand, we included all studies with quantitative performance measures. Importantly, we excluded participants who were not involved in organized sports teams or participating individually in an endurance sport. Burden et al⁵ found that those studies showing the largest effect on $\text{VO}_{2\text{max}}$ included participants with lower baseline $\text{VO}_{2\text{max}}$ values⁵; many of these studies were excluded from our review because we could not verify that their participants were involved in organized or endurance sports prior to starting the study.^{26,30}

Ferritin

There is a lack of standardization as to what constitutes a normal ferritin range in the current body of literature.^{11,19,20,27-29,35,36,39,47}

To make accurate and fair comparisons, we investigated whether there is a clinically applicable low ferritin cutoff that would support supplementation. A normal physiologic ferritin level varies between laboratories. However, levels less than 12 to 23 $\mu\text{g/L}$ are generally recognized as iron deficient, given a near absence of bone marrow iron stores.¹⁰ The sFer level at which investigators chose to start iron supplementation varied across the 12 studies examined in this review.^{4,11,19,20,27-29,35,36,39,47}

None of the studies in this review that used a ferritin level greater than 20 $\mu\text{g/L}$ as their cutoff demonstrated a significant improvement in performance following iron supplementation, whereas 6 of the 8 studies that used a cutoff level of 20 $\mu\text{g/L}$ or

less^{11,20,27,28,29,35,39,47} demonstrated an improvement in performance following iron supplementation.^{11,20,27,29,39,47} Therefore, it is reasonable to conclude that iron supplementation does not improve performance in athletes with sFer levels greater than 20 µg/L but that supplementation may play a role in improving performance in IDNA athletes with baseline sFer levels less than or equal to 20 µg/L.

Though many of the included studies measured other iron indices, only 1 study characterized athletes by stage 1 versus stage 2 iron deficiency.³⁵ Newhouse et al³⁵ made this distinction and indicated that 5 of the 19 participants in the iron treatment group had stage 2 iron deficiency. Notably, iron treatment did not improve performance (work capacity) even when participants with stage 2 iron deficiency were analyzed separately. Although this study did not demonstrate any significant difference in the performance of participants with stage 1 and stage 2 iron deficiency following iron supplementation, the hypothesis that athletes with more severe iron deficiency could benefit more from iron supplementation than those with less severe iron deficiency is plausible. Given the small sample size of the study by Newhouse et al,³⁵ future investigation into iron supplementation that distinguishes between the results of supplementation in stage 2 relative to stage 1 IDNA athletes is warranted.

Iron Supplementation Protocols

There is no consensus recommendation to guide the dosage, frequency, or delivery method and duration of iron supplementation in IDNA athletes. All of these parameters varied significantly among the review studies. Ferrous sulfate (20% elemental iron [the percentage absorbed by the gastrointestinal tract]) was the most common iron supplement employed in the studies reviewed. A typical recommended treatment for patients with IDA is 375 mg of ferrous sulfate taken twice daily; this is a total of 150 mg of elemental iron.²³ For reference, the recommended daily allowance of elemental iron for women (age 19-50 years) is 18 mg per day; the recommendation is only 8 mg per day for men.^{22,38} Based on the results of this review, there is no evidence to suggest that supplementation at or above the usually recommended daily allowance of iron improves performance in IDNA athletes. With regard to route of administration, none of the studies that used intramuscular or intravenous administration demonstrated improved performance.^{4,6,36} It is difficult to make any conclusive statements about the appropriate duration of iron supplementation for IDNA athletes. Of the studies that showed improved performance, the duration of treatment was highly variable with a very small sample size for each treatment duration; the studies that did not show improved performance had slightly more consistency, with 3 studies following participants for 8 weeks of treatment.

Limitations

The findings of this review are limited by the overall sample size as well as the demographics of participants. Although data from 12 studies were used, only 283 participants were included. In

addition, these studies included mainly young female participants, making it difficult to generalize results to all athletes.

CONCLUSION

Iron supplementation in IDNA athletes is a complex issue. The body of literature addressing whether iron supplementation improves performance in IDNA athletes is limited by small sample sizes, large discrepancies in performance measurement protocols, IDNA diagnostic protocols, and iron supplementation treatment protocols. The results of this review are equivocal and do not support the conclusion that iron supplementation consistently improves performance in endurance IDNA athletes. However, the studies that demonstrated performance benefits with iron supplementation in IDNA all used a ferritin supplementation threshold at or below 20 µg/L. There is no evidence to support the practice of iron supplementation to improve performance in IDNA athletes with sFer levels greater than 20 µg/L. Although the results of this review are equivocal regarding performance benefits following iron supplementation in IDNA athletes with ferritin levels less than or equal to 20 µg/L, we can conclude that iron supplementation in athletes with ferritin levels greater than 20 µg/L will not improve performance. This conclusion is significant in that a threshold of 20 µg/L can provide clinicians, athletes, and coaches with an evidence-based, clinically applicable threshold for iron supplementation in endurance athletes. Additionally, this review was not able to determine the optimal route or duration of treatment, since the reviewed studies used such a wide variety of supplementation regimens.

REFERENCES

1. Akabas SR, Dolins KR. Micronutrient requirements of physically active women: what can we learn from iron? *Am J Clin Nutr.* 2005;81:1246S-1251S.
2. Balaban EP. Sports anemia. *Clin Sports Med.* 1992;11:313-325.
3. Beard J, Tobin B. Iron status and exercise. *Am J Clin Nutr.* 2000;72:594S-597S.
4. Blee T, Goodman C, Dawson B, Stapf A. The effect of intramuscular iron injections on serum ferritin levels and physical performance in elite netballers. *J Sci Med Sport.* 1999;2:311-321.
5. Burden RJ, Morton K, Richards T, Whyte GP, Pedlar CR. Is iron treatment beneficial in iron deficient but non-anemic (IDNA) endurance athletes? A meta-analysis. *Br J Sports Med.* 2015;49:1389-1397.
6. Burden R, Pollock N, Whyte G, et al. Effect of intravenous iron on aerobic capacity and iron metabolism in elite athletes. *Med Sci Sports Exerc.* 2015;47:1399-1407.
7. Clarkson PM, Haymes EM. Exercise and mineral status of athletes: calcium, magnesium, phosphorus, and iron. *Med Sci Sports Exerc.* 1995;27:831-843.
8. Collins JF, Wessling-Resnick M, Knutson MD. Hepcidin regulation of iron transport. *J Nutr.* 2008;138:2284-2288.
9. Constantini N, Eliakim A, Zigel L, Yaaron M, Falk B. Iron status of highly active adolescents: evidence of depleted iron stores in gymnasts. *Int J Sport Nutr Exerc Metab.* 2000;10:62-70.
10. DellaValle DM. Iron supplementation for female athletes: effects on iron status and performance outcomes. *Curr Sports Med Rep.* 2013;12:234-239.
11. DellaValle DM, Haas JD. Iron supplementation improves energetic efficiency in iron-depleted female rowers. *Med Sci Sports Exerc.* 2014;46:1204-1215.
12. Dubnov G, Constantini N. Prevalence of iron depletion and anemia in top-level basketball players. *Int J Sport Nutr Exerc Metab.* 2004;14:30-37.
13. Dubnov G, Foldes A, Mann G, Magazani A, Siderer M, Constantini N. High prevalence of iron deficiency and anemia in female military recruits. *Mil Med.* 2006;171:866-869.
14. Ehn L, Carlmark B, Hoglund S. Iron status in athletes involved in intense physical activity. *Med Sci Sports Exerc.* 1980;12:61-64.

15. Fallon KE. Utility of hematological and iron-related screening in elite athletes. *Clin J Sport Med.* 2004;41:41-44.
16. Fischer RL, McMahon LF Jr, Ryan MJ, Larson D, Brand M. Gastrointestinal bleeding in competitive runners. *Dig Dis Sci.* 1986;31:1226-1228.
17. Fogelholm M. Indicators of vitamin and mineral status in athlete's blood cell count. *Int J Sport Nutr.* 1995;5:267-284.
18. Fogelholm M, Jaakkola L, Lampisjarvi T. Effects of iron supplementation in female athletes with low serum ferritin concentration. *Int J Sports Med.* 1992;13:158-162.
19. Fogelholm M, Rehunen S, Gref C, et al. Dietary intake and thiamin, iron, and zinc status in elite Nordic skiers during different training periods. *Int J Sport Nutr.* 1992;2:351-365.
20. Friedmann B, Weller E, Mairbaurl H, Bartsch P. Effects of iron repletion on blood volume and performance capacity in young athletes. *Med Sci Sports Exerc.* 2001;33:741-746.
21. Garza D, Shrier I, Kohl H, Ford P, Brown M, Matheson G. The clinical value of serum ferritin tests in endurance athletes. *Clin J Sports Med.* 1997;7:46-53.
22. Goodman C, Peeling P, Ranchordas M, Burke L, Stear S, Castell L. A to Z of nutritional supplements: dietary supplements, sports nutrition foods, ergogenic aids for health and performance—part 21. *Br J Sports Med.* 2011;45:677-679.
23. Harper J, Conrad M. Iron deficiency anemia. *Medscape.* <http://emedicine.medscape.com/article/202333-overview>. Accessed October 7, 2015.
24. Hassapidou M, Mantrantoni A. Dietary intakes of elite female athletes in Greece. *J Hum Nutr Diet.* 2001;14:391-396.
25. Haymes EM, Lamanca JL. Iron loss in runners during exercise: implications and recommendations. *Sports Med.* 1989;7:277-285.
26. Hinton PS, Giordano C, Brownlie T, Haas JD. Iron supplementation improved endurance after training in iron-depleted, nonanemic women. *J Appl Physiol (1985).* 2000;88:1103-1111.
27. Hinton P, Sinclair L. Iron supplementation maintains ventilatory threshold and improves energetic efficiency in iron-deficient nonanemic athletes. *Eur J Clin Nutr.* 2007;61:30-39.
28. Klingshirn L, Pate R, Bourque S, Davis J, Sargent R. Effect of iron supplementation on endurance capacity in iron-depleted female runners. *Med Sci Sports Exerc.* 1992;24:819-824.
29. LaManca J, Haymes E. Effects of iron repletion on $\text{VO}_{2\text{max}}$, endurance, and blood lactate in women. *Med Sci Sports Exerc.* 1993;25:1386-1392.
30. Magazanik A, Weinstein Y, Abarbanel J, et al. Effect of an iron supplement on body iron status and aerobic capacity of young training women. *Eur J Appl Physiol.* 1991;62:317-323.
31. Malczewska J, Blach W, Stupnicki R. The effects of physical exercise on the concentrations of ferritin and transferrin receptor in plasma of female judoists. *Int J Sport Med.* 2000;21:175-179.
32. McClung J, Karl P, Cable S, et al. Randomized, double-blind, placebo-controlled trial of iron supplementation in female soldiers during military training: effects on iron status, physical performance, and mood. *Am J Clin Nutr.* 2009;90:124-131.
33. McClung JP, Marchitelli LJ, Friedl KE, Young AJ. Prevalence of iron deficiency and iron deficiency anemia among three populations of female military personnel in the US Army. *J Am Coll Nutr.* 2006;25:64-69.
34. Newhouse IJ, Clement DB. Iron status in athletes: an update. *Sports Med.* 1988;5:337-352.
35. Newhouse IJ, Clement DB, Taunton JE, McKenzie DC. The effects of prelatent/latent iron deficiency on physical work capacity. *Med Sci Sports Exerc.* 1989;21:263-268.
36. Peeling P, Blee T, Goodman C, et al. Effect of iron injections on aerobic-exercise performance of iron-depleted female athletes. *Int J Sports Nutr Exerc Metab.* 2007;17:221-231.
37. Peeling P, Dawson B, Goodman C, Landers G, Trinder D. Athletic induced iron deficiency: new insights into the role of inflammation, cytokines and hormones. *Eur J Appl Physiol.* 2008;103:381-391.
38. Rodenberg RE, Gustafson S. Iron as an ergogenic aid: ironclad evidence? *Curr Sports Med Rep.* 2007;6:258-264.
39. Rowland T, Deisroth M, Green G, Kelleher J. The effect of iron therapy on the exercise capacity of nonanemic iron-deficient adolescent runners. *Am J Dis Child.* 1988;142:165-169.
40. Schumacher YO, Schmid A, Grathwohl D, Bültermann D, Berg A. Hematological indices and iron status in athletes of various sports and performances. *Med Sci Sports Exerc.* 2002;34:869-875.
41. Sinclair L, Hinton P. Prevalence of iron deficiency with and without anemia in recreationally active men and women. *J Am Diet Assoc.* 2005;105:975-978.
42. Smith JA. Exercise, training and red blood cell turnover. *Sports Med.* 1995;19:9-31.
43. Steen S, Mayer K, Brownell K, Wadden T. Dietary intake of female collegiate heavyweight rowers. *Int J Sport Nutr.* 1995;5:225-231.
44. Weaver CM, Rajaram S. Exercise and iron status. *J Nutr.* 1992;122(suppl 3):782-787.
45. Weight LM, Alexander D, Jacobs P. Strenuous exercise: analogous to the acute-phase response? *Clin Sci (Lond).* 1991;81:677-683.
46. Weight LM, Klein M, Noakes TD, Jacobs P. Sports anemia: a real or apparent phenomenon in endurance-trained athletes? *Int J Sports Med.* 1992;13:344-347.
47. Yoshida T, Udo M, Chida M, Ichioka M, Makiguchi K. Dietary iron supplement during severe physical training in competitive female distance runners. *Sports Med Train Rehabil.* 1990;1:279-285.
48. Zoller H, Vogel W. Iron supplementation in athletes—first do no harm. *Nutrition.* 2004;20:615-619.

For reprints and permission queries, please visit SAGE's Web site at <http://www.sagepub.com/journalsPermissions.nav>.